

LA-UR-12-23889

Approved for public release; distribution is unlimited.

Title: High Electric Fields in Cryogenic Liquids for EDM searches

Author(s): Ito, Takeyasu

Intended for: American Physical Society Division of Nuclear Physics Meeting 2012,
2012-10-24/2012-10-27 (Newport Beach, California, United States)



Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the Los Alamos National Security, LLC for the National Nuclear Security Administration of the U.S. Department of Energy under contract DE-AC52-06NA25396. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

DNP12-2012-000193

Abstract for an Invited Paper
for the DNP12 Meeting of
the American Physical Society

High Electric Fields in Cryogenic Liquids for EDM searches

TAKEYASU ITO, Los Alamos National Laboratory

Experiments to search for the neutron's electric dipole moment (EDM) using ultracold neutrons (UCNs) stored in superfluid liquid helium are under development. In these experiments, being able to achieve a strong and stable electric field in superfluid liquid helium in the region where UCNs are stored is of critically importance because in EDM searches in general the sensitivity depends linearly on the strength of the applied electric field. However, the phenomenon of electric breakdown in liquid helium is poorly understood, and as such major R&D efforts are under way to study it for these experiments. In this talk, the current status of such R&D efforts and the implications of the findings on EDM searches and on possible other applications will be discussed.